1/31/05-00287

Risk Management of Groundwater at Site 3, St. Juliens Creek Annex (SJCA), Chesapeake, Virginia

PREPARED FOR: SJCA Project Management Team

PREPARED BY: Kim Henderson/CH2M HILL

DATE: January 31,2005

This memorandum summarizes the potential risks identified and risk management conclusions for shallow and deep groundwater at Site 3. The potential risks to human and ecological receptors posed by media at Site 3 were identified in the *Final Remedial Investigation/Human Health Risk Assessment/Ecological Risk Assessnzeizt (RI/HHRA/ERA) Report for Sites 3*, 4, 5, and 6 (CH2M HILL, March 2003). The monitoring well locations at Site 3 are shown on Figure 1.

Shallow Groundwater (Columbia Aquifer)

Shallow groundwater is not considered a regional potable water source at or in the vicinity of SJCA due to its poor quality and low yield. Human health risks were evaluated for a construction worker scenario based on dermal contact and incidental ingestion only, and the results were below the Environmental Protection Agency's (EPA's) acceptable risk ranges (1x10⁻⁴ to 1x10⁻⁶ for carcinogens and Hazard Index of 1 for noncarcinogens).

Concentrations of one semivolatile organic compound (SVOC), bis(2-ethylhexyl)phthalate, and one inorganic analyte, thallium, exceeded their respective Maximum Contaminant Levels (MCLs) in shallow groundwater (Table 1). However;

- Bis(2-ethylhexyl)phthalate was detected in 1997 at a concentration (25 μg/L) above the MCL (6 μg/L) and the background upper tolerance limit (UTL) (1 μg/L) at only one monitoring well (SJS03-MW04S) and was not detected in subsequent monitoring rounds. Bis(2-ethylhexyl)phthalate is a known potential lab contaminant that likely reflects artifacts of the analytical process.
- Thallium was only detected at one monitoring well (SJS03-MW06S) and although the concentration (5.60 L μg/L) was above the MCL (2 μg/L), it was below the background UTL (7.6 μg/L). Additionally, a duplicate sample was collected at this location in which thallium was not detected. Therefore, thallium detected in shallow groundwater at Site 3 is not considered to be site-related.

Based on the discussion above, the shallow groundwater meets the MCLs in the latest round of sampling collected from Site 3 in 1999. Additionally, the SJCA Project Management Team (Navy, EPA, VDEQ) determined the previous MCL exceedances in shallow groundwater at Site 3 be anomalous and acceptable.

There is also the potential for groundwater from the Columbia Aquifer at Site 3 to discharge to sediment and surface water in the site drainages during high groundwater table

conditions. This exposure pathway was evaluated for potential ecological risks to aquatic life. Although several chemicals were identified as a potential concern in drainage surface water and sediment/soil that were also detected in groundwater, elevated concentrations of these chemicals were also detected in site soils. Based on the proximity of these media to one another, it is likely that surface soils represent the primary source of chemicals to the site drainages and this pathway was mitigated by the removal of waste, soil, and upland drainage sediment/soil at Site 3. Thus, groundwater does not appear to be a significant transport route from the site to the drainages.


Deep Groundwater (Yorktown Aquifer)

In the area of SJCA, the deep groundwater is not used as a potable water supply. Area residences are only located upgradient of Site 3 and are supplied potable water by the City of Chesapeake. Deep aquifer groundwater is currently only used for irrigation upgradient of SJCA. Although future residential development of the site is improbable, deep groundwater was evaluated under the future residential scenario. Table 2 shows the deep groundwater data collected at Site 3.

Potential noncarcinogenic hazards (based on ingestion of arsenic, manganese, and thallium) and carcinogenic hazards (based on contact with arsenic) were identified to current and future adult and child residents who use the deep groundwater beneath Site 3 as a potable water supply. However;

- Arsenic was only detected in one unfiltered sample in the upgradient monitoring well (SJS03-MW01D) at an estimated concentration (4.20 J μ g/L) below the MCL (10 μ g/L).
- Manganese concentrations (ranging from 45.6 μ g/L to 612 μ g/L) were low and below the tap water risk-based concentration (RBC) of 730 μ g/L.
- Thallium was only detected in a 1997 sample at an estimated concentration (2.7 J μ g/L) slightly above the MCL (2 μ g/L) from the upgradient monitoring well (SJS03-MW01D) and not detected in the subsequent monitoring events.

Based on the discussion above, the deep groundwater meets the MCLs in the latest round of sampling collected from Site 3 in 1999. Additionally, the SJCA Project Management Team (Navy, EPA, VDEQ) determined the deep groundwater risks identified at Site 3 to be acceptable for all pathways and receptors.



	Screenk	o Criteria																		
Station ID	RRC-Tax	ORGIT MCL Back SJS03-MW01S			\$J\$G3-MW028					\$JS03 MW038			\$J\$03-NW04\$				SJS02 MY COSLS SJS02 MY		3-MW06S	
Sample ID	(10/02)	Ground	ground UTLs	SJS03-GW1S-001	SJS03-GW15-002	SJS03-GW1S-003	\$J\$03-GW2S-001	SJS03-GW2S-002	SJ\$03-GW2S-003	SJS03-GW3S-001	SJS03-GW3S-001P	SJS03-GW3S-002	\$JS03-GW3S-003	SJS03-GW4S-001		SJS03-GW4S-002P	SJS03-GW49-003	SJ503-GW55-001	SJ\$63-GW6\$-001	SJS03-GW6S-001P
Sample Date		water	Oifs	07/17/97	11/04/97	05/23/99	07/17/97	11/04/97	06/19/99	07/21/97	07/21/97	11/04/97	05/19/99	07/17/97	11/04/97	11/04/97	06/19/99	06/19/92	05/19/99	05/19/99
Chemical Name																		1		
VOCs (UG/L)																	ļ	 		
Acetone	610		30.9	24	NA NA	NA.	8	5 U	NA	50	5 J	10	NA	50	6 U	60	NA.	144	17	17 3
Ethylberzene	3.3	700	0.6	0.6.7	10	10	10	10	10	10	10		10	10		10		10	10	10
Totuene	750		0.2	02.J	1.0	10	10	10	10	1.0	10		10	10	10	10			1 0	10
Xylene, total	12,600	10,000		1	10	10	1 U	10	11/	10	1 0	1 U	1.0	10	10	10	10	10	. 10	10
SVOCs (UG/L)		 -								ļ				<u> </u>					l	
2-Methylnaphthalene	120		21	1.7	10 (3	11 U	10 U	10 Ú	11 U	16.0	10 U	10 U	11.0	10 U	16 U	10 U	110	110	10 U	11.0
d-Methylphenol	180		NC)	10 U	10 U	11 U	10 U	10 ()	11 U	10 U	10 U	10 U	11 0	10 U	10 U	10 0		110	2.1	
Aceraphthene	370	-	18	15	4 3	3.3	10 U	10 U	11 U	2 J	2 J	7 J	3.3	10 U	10 U	10 U		11 U	10 U	11 0
Cartiazole	3.3		2	2 J	10 U	11 0	10 U	16 U	11 U	10 U	10 U		11 U	10 U	10 Ų	10 U		11 U	10 U	11.0
Oitenzoluran	1,500			8 J	1.3	11 0	10 U	10 U	51.0	10 U	10 U	10 U	11 U	10 U	10 U	10 U		11 U	10 U	11.0
Fluoranthene Fluorene	1,500			10 (1	1 J 10 U	11 U	10 U	10 U	11 0	10 U	10 U	10 U	11 U	10 U	10 U	10 U	11 0	11 U	10 U	11 U
Naptshalene	6.5		19	19	10 U	11 U	10 U	10 U	11 0	10 U	10 U	10 U	11 0	10 U	10 U	10 0		11 0	10 U	11 0
Phenarthrene	180		B G	6 J	10 U	11 U	10 U	10 U	11 U	10 U	10 U	10 U	11 U	10 U	10 U	10 U	11 U	11 0	10 U	11.0
Phenol	11,000	-	ΝÜ	10 U	10 U	11 0	10 U	10 U	11 U	10 U	10 U	10 U	11 U	10 U	10 U	10 U	11 U	11.0	2.1	2 J
tis(2-Eëryiheryi)ptehalate	4.8	6	1	10 U	10 U	1 J	10 U	10 U	11 U	10 U	2 J	10 U	11'U	25	14 B	7 B	11 U	11 U	10 U	11 U
o	ļ	\vdash																		
PessPCDs (UGA.) Heptachlor	0.015	0.4	Pit.	0.06.0	0.05 U	0.0520 U	0.06 U	0 05 U	0.0240 J	0.05 U	0.05 U	006 U	0.0670 U	0 0 S U	9 05 U	0 05 U	0 0570 U	0.0520 U	0 0620 U	0 0520 U
фили		<u>V.</u>	- "			000200	0.00 0	0000	W. 1470 J	0.000	0030	0.60	0.0870 0	0050	0000	0050	005/00	0.0520.0	0.6200	0.0520.0
Fotal Metals (UG/L)								7						1				1		
Alumnum	37,000		1,710	4,100 K	1,710	395	19,200 K	7,660	6,850	4,460 K	8,270 K	38.1 U	186 B	9,170 K	56.8 J	3810	247 B	621	92.7 B	55 5 B
Artimony	15		23	228	2.3 J	2.70 U	2 U	1.7 U	270 U	2.0	2.0		2.70 U	2 U	1.7 U	1.7 U	270 U	2.70 ∪	2.70 U	270 U
Amenic Barium	2,600	2,000	77.4	4 Å J 176 J	32 U	3,46 J 23.1 J	3 U 76 G J	3.4 J 24.7 J	4.70 J 26 6 J	3 U 43 J	3 U 476.1		2 U	30	320	320	2 U	610 J	6 80 J	4.10.3
Seryllium	73		1.4	14.3	0.76 J	0.100 U	10	24.13	20.63	10	1 0	16 1 J 0 96 J	126 J 0 100 U	61.4 J	207 J 0.95 J	20 7 J 1.2 J	213J	121 J 0 100 U	143 J	147 J 0 100 U
Cadmium	18	- 6	074	0.5 U	0.4 U	0.420 J	0.6 U	04 U	1.10 J	060	0.5 U	04 U	0300 U	060	04 U	04 U	0300 ∪	0.350 J	0490J	0.360 J
Calcium		-	531,000	100,000	92,000	141,000	365,000	488,000	529,000	89,700	95,900	367,000	201,000	178,000	130,000	129,000	156,000	194,000	226,000	212,000
Chronium	110	100	3.2	16.1 B	12 B	3 20 J	56.7	4.6 U	3 10 J	13.8 B	183 B	4 6 U	1,10 U	35.2	46 U	46 U	1.10 U	1,60 J	1,10 U	1 16 U
Cebalt	730 1,500	1,300	158	8.5 J 7.4 B	63 J	2.30 J 6.20 B	16 J 159 B	50.7 5.8 U	16.6 J	8 U 68 B	8 U	6 U	0.5 U 4.10 B	80	38.5 J 58 U	36.8 J 58 U	6,10 J 340 B	6.10 J 10 6 B	0 690 B	1.20 B 4 60 J
Copper	11,000	1,30.	107,000	31,900	26,200	34,600	33,900	19.400	35 300	32,900	35 000	45.000	24 600	16,900	20,100	20.000	20,600	26,100 10 6 B	25400	16,100
Lead	15	15	3.5	24 B	3.5	2 30 J	9	130	190 J	26 J	4.3	17 J	140 J	5.7	143	130	10	10	10	10
Magnesium			295,000	115,000	34,600	31,600	179,000	168,000	179,000	75,300	77,000	33,600	15,000	154,000	48,900	46,700	54,200	128,000	210,000	300,000
Manganese	736		13,700	2,960	1,720	1,770	2,990	6,330	5,410	830	ESO	2,960	1,600	1,030	1,320	1,320	1,440	2,610	1,535	1,440
Nickel Potavskim	730		26 1 85,400	7 U 37,300	14 5 J 26,600	4.3	25.1 J	84.8	21 J	7 U	7 0	63 U	0.900 U	7 U	46.1	43.1	5 J	7.90 J	1,70 J	160 J
Seleraum	160	7/	10,400 NC	37,300	26,000 3 3 U	27,100 2.60 U	61,100 3 U	25,100 3.3 K	35,400 2.60 U	30,200 3 U	31,000 3 U	23,000	13,900 2.60 U	57,000 3.U	28,200	27,600 33 U	31,000 2 60 U	33,600 2 60 M	83,400 3.20 B	81,300 2 60 U
Silver	180		1.9	1.9 J	1.1 U	0 900 U	10	1.1 U	0.900 U	1.4 8	10		0,900 U	23B	17 J	7.7		0,900 U	0.900 U	0,900 U
Sodium			810,000	810,000	122,000	132,000	612,000	293,000	376,000	470,000	468,000	39,400	20,800	799,000	155,000	154,000	182,000	1,080,000	1,660,000	1,740,000
Thallium	2.6	- 2	7.6	4.6 9	3.1 B	3.20 U	2 U	1.5 U	3 20 U.	280	24 8	16 U	3 20 UL	2 U	15 U	1.5 U	3.20 UL	3 20 UL	5 (O L	3.20 U.
Vanadium V	260		13.7	13 7 J	12.2 J	1.5 J	43 J	68 U	0.600 U	18.5 J	22.7 J	680	0,600 U	15.5 J	68 U	68∪	0,600 U	0810 J	0.650 J	0.600 U
zine	11,000		241	35.3 8	65 2 B	241	90	397	46.9	20.8 B	22 8	15 f B	640.3	36 4 B	327	327	28.6	138 J	1,90 U	190 U
Dissolved Metals (UGAL)																	l	1		
Alummum	37,000	-	399	44 U	64.2 8	67.1 Đ	44 U	807	116 B	149 B	164 8	38.1 U	38 2 U	44 U	38 1 U	38,1 U	161 8	70.8 B	60 1 B	44.1 8
Antimony	15	- 6	38	2 U	3.8 J	2.70 U	2 U	1,7 U	270 U	2 U	2 U	1,7 U	2.70 U	2 U	1,7 U	1,7 U	270 U	2.70 U	2.70 U	2 70 U
Arseric	0.045	10	24	3 U	3.2 U	2400	3.0	3.2 ∪	2 30 J	3 U	3 U		2 ()	3 U	32 U	32 U	2 U	6.3	L 3.8	5 10 6
Sarium Beryllium	2,600	2,000	93.3	295 1 U	26 J 0 58 U	21.4 J	43.4 J	25 J	24 J 0 100 U	37.6 J	44.3 J	14.9 J 0.58 U	10.5 J 0.100 U	38 J	23 6 J	22.1 J 1.2 J	21.3 J	0,100 U	175 J 0,100 U	161 J 0,100 U
Cadmium	18		0.31	050	0.8 U	0.330 J	1.U 0.5.U	049 J	0.100 U 0.540 J	060	05 U		0 100 U	05 U	9.2 J	1.2 J	0.510 J	0.100 U	0.100 U	0.560 J
Calcium	- "	<u> </u>	464,000	148,000	93,500	132,000	348,000	524,000	513,000	87,500	98,200	340,000	202,000	178,000	139,000	133,000	150,000	154,000	223,000	198,000
Chromium	110	100	58	7 U	58J	1.60 J	7.0	4.6 U	1.10 U	928	12.5 B	4.6 U	1.10 U	7 U	46U	4.6 U	1.10 U	140 J	1 80 J	2.5 J
Coball	730		15	8 U	7.7 J	2.10 J	12.2 J	63	16.1 J	8.0	80		0.5 U	8.0	429 J	41.8 J	? J	4.30 J	050	0.5 U
iron	11,000		94,000	74,500	24,900	31,900	10,900	20,000	34,000	30,000	33,600	42.200	24,100	6,990	21,400	20,766	20,100	38,000	42,300	42,100
Lead Magnesium	15	16	256,000	1 U 194,000	1.9 B	1 U 32,900	1 U 170,000	1,4 J 183,000	1 U	77,800	1 U 86,890	1.3 U 31,900	1 U 15,000	1 U 150,000	1.3 U 52,500	1,3 U 50,600	1 U 51,700	1 U 131,000	1 30 J 323,000	1 U 799,000
Manganese	730		11,800	3,070	1,760	1,790	2,770	6,710	6,180	800	80,800 911	2700	1,120	1,020	1,420	1,370	1,400	2,930	1,760	1,840
Nickel	730		13.2	7 U	83B	3 10 J	7 U	89.1	77.1 J	10	7 U	63 U	0 900 U	7 U	50.5	45,7	75 J	7.29 J	1.5 J	2898
Potassium			73,300	38,900	27,000	28,400	57,200	27,100	33,100	30,800	34,600	21,400	13,500	56,000	30,300	29,000	29,900	28,600	85,400	75,000
Selenum	180	50	NO	3 U	3.3 U	2.60 U	3 U	1.3 3	260 ∪	3 U	3.0		2.60 U	3 ∪	33 U	33 U	4.5 J	260 U	2 60 U	3 90 B
Selver Sectorn	180		24	24 J	1.5 8	0 900 U	1.0	1.10	0 900 U	12 B	10	2 8	0,900 U	10	1 6 B 168,000	2 7 B 161,000	0 900 U 172,000	0 900 U	0 900 U 2,120,000	0 990 U
Sedeum Vanadeum	260		682,000 7 1	1,530,000	125,600 7.1 J	141,000 1,20 J	537,000 9 U	307,000 6 E U	362,000 0 600 U	525,000	557,000 9.U	38,700 68 U	20,200 0.600 U	793,000 S U	168,000 6 8 U	161,000 68 U	0 600 U	0.600 U	2,120,000 0,600 U	0,000 U
Zinc	11,000		109	224 8	49.7 B	105 J	27.3 B	497	36	20 5 B	1178	24.4.8	137 J	13 5 6	325	111	1093	218	1,90 U	190 U
	1.,430													Q.,,,,,,,,,,			 			

Exceeds RDC/Lip
Exceeds RDC/Lip
Exceeds RdC/Groundwater
Exceeds Rackground UTL
B. Analyte not detected above associated blank
J. Reported value in estimated
K. Reported value may be bis sed high.

L - Reported value may be biased low

1Nx - 1804 Analyzed

1Dx - Compounds were analyzed but not deserted duming Background Investigation

185 - Compounds not analyzed during Background Investigation

P. Duplicate sample

- Analyze not deserted

Table 2 Deep Groundwater Exceedances of Screening Criteria at Site 3

St. Juliens Creek Annex

Chesa	peake.	. Vir	ninia	a

	Screening	Criteria		Chesapeake, Virginia									
Station ID	RBC-Tap	MCL-				SJS03-MW01D	SJS03-MW02D						
Sample ID	(10/02)	Ground	BACKGROUND1		SJS03-GW1D-001	SJS03-GW1D-002	SJS03-GW1D-003	SJS03-GW2D-001	SJS03-GW2D-002	SJS03-GW2D-003			
Sample Date	(10/02)	water	Mean	Max	07/24/97	11/04/97	05/19/99	07/24/97	11/04/97	05/19/99			
Chemical Name													
VOCs (UG/L)													
Carbon disulfide	1,000		ND	ND	1 U	0.3 J	0.200 J	1 U	1 U	7.60			
Chloroform	0.15	80	ND	ND	0.4 J	1 U	1 U	0.3 J	1 U	1 U			
SVOCs (UG/L)													
bis(2-Ethylhexyl)phthalate	4.8	- 6	4.0	1	10 U	3 J	12 U	10 U	10 U	1 J			
Pest/PCBs (UG/L)	 								***************************************				
No Detections			ļ										

Total Metals (UG/L)	0.045				0.11		1001	0.11	0011	0.11			
Arsenic Barium	0.045 2,600	2,000	ND 24.7	ND 32.7	3 U 15.8 J	3.2 U 21.7 J	4.20 J 30,7 J	3 U 46.6 J	3.2 U 49.1 J	2 U 60.2 J			
Calcium	2,000	2,000	61,933	75,200	49,300 J	49,500	61,700	103,000 J	95,200	110,000			
Chromium	110	100	01,933 ND	75,200 ND	49,300 J 7 U	49,500 4.6 U	1,40 J	7 U	95,200 4,6 U	2.40 J			
Iron	11,000	100	907.7	1.670	788 J	170 B	476	487 J	272	541			
Lead	15	15	907.7 ND	1,570 ND	1 UL	1,8 J	1 J	2.1 B	1.3 U	1 U			
Magnesium	i	- 10	6,630	12,000	14,400 J	12,700	16,600	39,000 J	37,100	48,000			
Manganese	730		177.7	226	91.3 L	45.6	110	291 L	354	612			
Nickel	730		0.7	1.1	8.1 J	6.3 U	0.900 U	7 U	6.3 U	1 J			
Potassium			4,817	8,490	15,100	15,400	16,600	17,700	16,000	17,600			
Silver	180		ND	ND	1 U	1.1 U	0.900 U	1 U	1.2 J	0.900 U			
Sodium			30,233	39,900	40,600 J	43,300	50,600	37,300 J	34,500	41,000			
Thallium	2.6	2	ND	ND	2.7 J	2.5 B	3.20 UL	2 U	1.5 B	3.20 UL			
Zinc	11.000		3.5	4.8	6.4 B	10.9 B	3.80 J	7.1 B	9 B	16.6 J			
					annual control and a series and								
Dissolved Metals (UG/L)													
Barium	2,600	2,000	23.0	31.2	14.1 8	20.8 J	23.6 J	42.8 J	53.7 J	58.7 J			
Calcium			57,667	77,700	53,800 J	51,100	63,700	98,400 J	103,000	105,000			
Iron	11,000		788.7	1,360	180 B	130 B	301	156 B	276	464			
Magnesium			5,967	10,000	18,700 J	13,200	18,000	38,200 J	40,900	46,300			
Manganese	730		161.3	188	101 J	46.4	123	284 J	385	583			
Nickel	730		1.3	2.3	7 U	6.3 U	0.980 J	7 U	6.3 U	0.900 U			
Potassium			4,407	7,200	12,600	15,100	17,900	16,600	17,600	18,100			
Sodium			28,400	37,900	44,100 J	44,800	52,900	35,200 J	37,800	39,600			
Thallium	2.6	2	ND	ND	2.1 J	1.5 U	3.20 U	2 U	1.9 B	3.20 U			
Zinc	11,000		6.1	10.5	5 U	5.7 B	4.40 J	7.2 B	18.5 B	10.5 J			

Notes:

Exceeds RBC-Tap

Exceeds MCL-Groundwater

NA - Not Analyzed

B - Analyte not detected above associated blank

J - Reported value is estimated

L - Reported value may be biased low

NO -Compounds were analyzed but not detected during Background Investigation

U Analyte not detected

Potential human health risk drivers

¹ Background UTLs have not been established for deep gorundwater at SJCA. The concentrations shown are mean and maximum detected concentrations from sampling conducted at 3 deep monitoring wells in 1999.